

최댓값과 최솟값

(Maximum and Minimum Values)

Maximum and Minimum Values

▶ Start

▶ End

▶ Start

▶ End

Definition

▶ Start

▶ End

Definition

M :

▶ Start

▶ End

Definition

M : absolut maximum value

▶ Start

▶ End

Definition

M : absolut maximum value of f

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$\exists c \in D$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$\exists c \in D$ s.t. $M = f(c)$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$\exists c \in D$ s.t. $M = f(c) \wedge$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$\exists c \in D$ s.t. $M = f(c) \wedge [x \in D$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m :

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c)$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M :

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D$$

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c)$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow$$

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

▶ Start

▶ End

Definition

M : absolut maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolut minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m :

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value of f

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value of f on D

$$\exists c \in D$$

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0$$

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } m = f(c)$$

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } m = f(c) \wedge$$

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } m = f(c) \wedge [|x - c| < \varepsilon$$

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } m = f(c) \wedge [|x - c| < \varepsilon \Rightarrow$$

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } m = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \leq f(x)]$$

▶ Start

▶ End

Definition

M : absolute maximum value of f on D

$$\exists c \in D \text{ s.t. } M = f(c) \wedge [x \in D \Rightarrow f(c) \geq f(x)]$$

m : absolute minimum value of f on D

$$\exists c \in D \text{ s.t. } m = f(c) \wedge [x \in D \Rightarrow f(c) \leq f(x)]$$

M : local maximum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } M = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \geq f(x)]$$

m : local minimum value of f on D

$$\exists c \in D \wedge \exists \varepsilon > 0 \text{ s.t. } m = f(c) \wedge [|x - c| < \varepsilon \Rightarrow f(c) \leq f(x)]$$

Github:

<https://min7014.github.io/math20240220001.html>

Click or paste URL into the URL search bar,
and you can see a picture moving.